

Dependence of the Fractional Quantum Hall Edge Critical Exponent on the Range of Interaction

Recent experiments on external electron tunneling into an edge of a fractional quantum Hall (FQH) system find a striking current-voltage power law behavior $I \propto V^\alpha$, with exponent $\alpha \approx 2.7$ on the $f = \frac{1}{3}$ FQH plateau [1,2]. Such power law I - V characteristic with $\alpha = 3$ was predicted for electron tunneling into an edge channel at the boundary of the $f = \frac{1}{3}$ FQH system: the low-energy dynamics is effectively 1D, and field-theoretic descriptions of edge channels as chiral Luttinger liquids have been developed [3]. The equality $\alpha = 2i + 1$ for a FQH state at $f = \frac{1}{2i+1}$ ($i = 1, 2, 3$) has been demonstrated theoretically in the disk geometry for the Laughlin wave function Ψ_L , which is known to be the exact ground state for certain short-range interactions. The experiments consistently obtain values of $\alpha < 2i + 1$, however [2].

Here we report results of a large numerical study of the microscopic structure of the FQH edge at $f = \frac{1}{3}$. To this end, we diagonalize the interaction Hamiltonian in the disk geometry for up to $N = 12$ spin-polarized electrons restricted to lowest Landau level. We construct numerically [4] the Laughlin Ψ_L and “Coulomb” Ψ_C wave functions as the ground states of the short-range and Coulomb Hamiltonians, respectively, for the total angular momentum $M = \frac{3}{2}N(N-1)$, which gives filling $f = \frac{1}{3}$ in the thermodynamic limit $N \rightarrow \infty$. In particular, we have obtained occupation numbers $\rho(m)$ of the angular momentum basis orbitals

$$\psi_m(r, \theta) = (2\pi 2^m m!)^{-1/2} r^m \exp(im\theta - r^2/4), \quad (1)$$

where radius r is in units of magnetic length $\ell = \sqrt{\hbar/eB}$. The Hilbert space is restricted by consideration of orbitals with angular momentum $m \leq m_{\max}$ only. For the Haldane V_1 short-range interaction, the Laughlin $\rho_L(m)$ with $m > m_{\max}^L = 3(N-1)$ vanish identically; for Coulomb interaction good convergence of $\rho_C(m)$ is obtained by $m_{\max} = m_{\max}^L + 5$ [6]. For example, for $N = 12$, the largest $f = \frac{1}{3}$ FQH system studied, $M = 198$ and the size of the Hilbert space is 15,293,119 for $m_{\max} = 35$. Details of this study will be published elsewhere [5].

As has been demonstrated by Wen [3], the critical exponent α is equal to the ratio of the occupation numbers

$$\alpha = \rho(m_{\max}^L - 1)/\rho(m_{\max}^L) \quad (2)$$

for the Laughlin state Ψ_L on the disk. Wen has also argued that this relationship must hold for any interaction, so long as the FQH state at the same filling f exists, and is not unique for the Laughlin wave function. In Fig. 1 we present the ratio of the occupation numbers for both Ψ_L (short-range interaction) and Ψ_C (true Coulomb interaction) for $N = 3$ to 12. As expected, we obtain $\alpha_L = 3$ to machine accuracy for Ψ_L . For Coulomb-interacting




FIG. 1. The ratio of the angular momentum occupation numbers $\rho(m)$ for N interacting electrons on the disk. Shown are the Laughlin state ρ_L for a short-range interaction, and the exact ground state for the Coulomb interaction ρ_C .

electrons, the ratio is always less than 3, and an extrapolation to the thermodynamic limit gives $\alpha_C = 2.62$. While we do not know whether the extrapolation shown in Fig. 1 holds for $N > 12$, certain other systematic behavior present in the numerical data [5] allows us to project that in the $N \rightarrow \infty$ limit $2.58 \leq \alpha_C \leq 2.75$, and is definitely less than 3.

Thus we propose that the deviation of the experimental α from the predicted α_L values is not an artefact, and is not due to corrections such as finite bulk diagonal conductivity, disorder, or a variation of the electron density in the sample. Rather, we propose, the effect has a fundamental origin: the value of the critical I - V exponent is not universal, but depends on the range of particular interaction.

This work was supported in part by the NSF.

V. J. Goldman and E. V. Tsiper

Department of Physics, State University of New York
Stony Brook, NY 11794-3800

January 11, 2001

- [1] A. M. Chang, L. N. Pfeiffer, and K. W. West, Phys. Rev. Lett. **77**, 2538 (1996); M. Grayson *et al.*, Phys. Rev. Lett. **80**, 1062 (1998).
- [2] A. M. Chang *et al.*, Phys. Rev. Lett. **86**, 143 (2001).
- [3] X. G. Wen, Phys. Rev. B **41**, 12 838 (1990); Intl. J. Mod. Phys. B **6**, 1711 (1992).
- [4] E. V. Tsiper and V. J. Goldman, cond-mat/0101151.
- [5] E. V. Tsiper and V. J. Goldman, to be published.
- [6] The sum of $\rho_C(m)$ with $m > m_{\max}^L$ is less than $0.01/N$. This means that fixing total angular momentum $M = \frac{3}{2}N(N-1)$ selects the $f = \frac{1}{3}$ state for Coulomb interaction too: it fixes *average* density on the disk $\langle \rho_C \rangle \approx \langle \rho_L \rangle$ for $r < r_{\text{edge}}$ to better than 10^{-3} for any N .